

Group Theory Approach to Band Structure: Scarf and Lamé Hamiltonians

Hui Li and Dimitri Kusnezov *

Center for Theoretical Physics, Sloane Physics Laboratory, Yale University, New Haven, CT 06520-8120

(November 30, 1998)

The group theoretical treatment of bound and scattering state problems is extended to include band structure. We show that one can realize Hamiltonians with periodic potentials as dynamical symmetries, where representation theory provides analytic solutions, or which can be treated with more general spectrum generating algebraic methods. We find dynamical symmetries for which we derive the transfer matrices and dispersion relations. Both compact and non-compact groups are found to play a role.

The application of group theoretical techniques to physical problems has a long and fruitful history. From the early works on the low-lying spectrum of hadrons [1], to the more recent studies of nuclear [2] and molecular [3] many-body problems, the theory of Lie groups and algebras has become important not only in explaining the behavior of various physical systems, but also in constructing new physical theories. Problems of interest can usually be approached in this manner when a spectrum generating algebra (SGA) can be identified. A SGA exists when the Hamiltonian H can be expressed in terms of generators of an algebra. As a consequence, the solution of the Schrödinger equation then becomes an algebraic problem, which can be attacked using the tools of group theory. In addition to providing a general approach to solving the problem of interest, the SGA also provides as special cases, exactly solvable limits of the models. These are known as dynamical symmetries, and can often be found by suitable choices of the parameters of a model. Typically one has more forms of the Hamiltonian which are exactly solvable, than dynamical symmetries predicted from group theory [4].

The dynamical symmetry problems up to now have focussed essentially on the description of bound and scattering states. By mapping the quantum mechanical problem onto an algebraic structure, representation theory immediately provides the excitation energies and eigenstates. However, there is one missing category in the group theoretical description of quantum systems, namely, Hamiltonians with periodic potentials. In this case, the energy spectrum is characterized by energy bands and gaps. We extend the group theoretical framework by showing that dynamical symmetry and SGA techniques can be applied to such problems, where dispersion relations, transfer matrices and energy bands can be understood in the context of representation theory.

Consider Hamiltonians in one dimension of the form:

$$H = -\frac{\hbar^2}{2M} \frac{d^2}{d\rho^2} + V(\rho). \quad (1)$$

For the Pöschl-Teller potential, $V(\rho) \propto 1/\cosh^2(\rho/\rho_0)$, it is well known that bound states can be expressed as discrete representations of $SU(2)$, and scattering states by the principal series of $SU(1,1)$, for which $j = -1/2 + ik$, $k > 0$ [5–7]. In passing from $SU(2)$ (where $[J_z, J_{\pm}] = \pm J_{\pm}$ and $[J_+, J_-] = 2J_z$) to $SU(1,1)$ (obtained by changing the sign in $[J_+, J_-] = -2J_z$), the Casimir invariant, J^2 changes from $J^2 = J_x^2 + J_y^2 + J_z^2$ (sphere) to $J^2 = J_z^2 - J_x^2 - J_y^2$ (hyperboloid), and the representation quantum numbers j, m fall into three well known series: the principal, discrete and complementary. However, if one has a periodic potential $V(\rho) = V(\rho + a)$, for some constant a , it is not at all clear how this fits into the scheme provided by representation theory. In this case we would like to derive the transfer matrices and dispersion relations.

When one considers the eigenstates in an energy band, Bloch's theorem states that it must have the form $\Psi_k(\rho) = \phi_k(\rho) \exp(ik\rho)$, where $\phi_k(\rho + a) = \phi_k(\rho)$ has the periodicity of the potential. Group theory typically provides single-valued wavefunctions, suitable for band edges. In the $SU(1,1)$ dynamical symmetries we construct, we will see that wavefunctions can be obtained by (a) using multiple-valued projective representations (j, m) (labeled by J^2 and J_z and associated with equivalence classes of vectors defined up to a phase, which is natural in Hilbert space), and (b) by using a basis $|j, \lambda\rangle$ labeled by J^2 and a non-compact generator of $SU(1,1)$ such as J_x . Several surprising aspects will emerge, such as the use of the complementary series and the need for non-unitary representations.

In order to see how dynamical symmetry methods and representation theory might apply to Hamiltonian with band structure, we must first construct a dynamical symmetry for a periodic potential. It is possible to do so with the Scarf potential [8]: $V(\rho) \propto 1/\sin^2(\rho/\rho_0)$. In dimensionless variables:

$$H_{Sc} \Psi(x) = \left[-\frac{d^2}{dx^2} + \frac{A}{\sin^2 x} \right] \Psi(x) = \mathcal{E} \Psi(x). \quad (2)$$

(The energy is now $E = (\hbar^2/2M\rho_0^2)\mathcal{E}$.) This potential is known to have band structure for $-\frac{1}{4} < A < 0$ [8]. By suitable choices of generators, we can realize H_{Sc} as a Casimir invariant. There are different ways to do this,

but not all are fruitful. For instance, in $SU(2)$ we obtain $A = m^2 - 1/4$, and $\mathcal{E} = (j+1/2)^2$. Here $H_{Sc} = \mathbf{J}^2 + 1/4$ is simply related to the Casimir invariant $\mathbf{J}^2$ of $SU(2)$. This realization is unsuitable since (i) the discrete representations give only periodic solutions and (ii) the coupling A is never in the range of physical interest.

If we choose the generators of $SU(1,1)$ in the form $J_{\pm} = e^{\pm i\phi}(\mp \sin \theta \partial_{\theta} + i \cos \theta \partial_{\phi})$, and $J_z = -i\partial_{\phi}$, we can realize H_{Sc} with $A = j(j+1)$ and $\mathcal{E} = m^2$:

$$H_{Sc}\Psi_{jm} = J_z^2\Psi_{jm} = m^2\Psi_{jm}. \quad (3)$$

This is more natural, since a given strength of the potential corresponds to a fixed representation j , and the eigenvalue is related to the projection m . (We have replaced θ by x in relating Eq. 3 for Eq. 2). To describe band spectra in $SU(1,1)$, we pass to the projective unitary representations [9], which have three classes: the discrete series $D_j^{\pm}$, with $j < 0$, the principal series where $j = -1/2 + i\sigma$, ($\sigma > 0$), and the complementary series with $-1/2 < j < 0$. While the discrete and principal series of $SU(1,1)$ has found application to problems such as the Pöschl-Teller potential discussed above, the remaining complementary series has found little or no application in physics. Surprisingly, this series is precisely what is needed for band structure, and is essential to complete the dynamical symmetry approach to quantum systems.

The complementary series is shown in Fig. 1 (top) and corresponds to potentials with $-1/4 < A < 0$, exactly the region of interest. The unitary representations are labeled by two quantum numbers j, m , with $m = m_0 \pm n$ ($n = 0, 1, \dots$), where $0 \leq m_0 < 1$ and $m_0(m_0 - 1) > j(j+1) \geq -1/4$. In the figure, we plot a generic line $j(j+1) = \text{const}$ (dots) together with the parabola $m_0(m_0 - 1)$ (solid). We also show the periodic repetition of this parabola given by $(m \pm n)(m \pm n - 1)$ (dashes). The values of j, m which correspond to unitary representations are then found to be $n-j \leq m \leq j+n+1$ as well as $-(j+n+1) \leq m \leq -(n-j)$. The relation of these representations to the band structure is shown in Fig. 1(bottom). There we plot the eigenvalue m^2 as a function of j , related to the strength of the potential through $A = j(j+1)$. The range of m corresponding to unitary representations give the *energy gaps*. The *non-unitary* representations correspond to the *energy bands* (shaded region). We know that the non-unitary representations must correspond to the energy bands from an analysis of the $j \rightarrow 0$ limit. Here the strength of the potential $A = 0$, and we have free motion, where all values of m must be allowed, which from Fig. 1(top) corresponds to the non-unitary representations.

The discrete (projective) representations are readily understood as well. The series $D_j^{\pm}$ provides the lower band edges, as shown by the boxes in Fig. 1(bottom) for a particular choice of potential given by j (dashes). (Both series $\pm$ are degenerate and give the same spectrum.) The upper band edges are obtained by realizing

that the strength $A = j(j+1)$ of the potential is invariant under the transformation $j \rightarrow -(j+1)$. The discrete series $D_{-1-j}^{\pm}$ then corresponds to the upper band edges (circles). The principal series has a potential with strength $A < -1/4$, for which the Hamiltonian is no longer self-adjoint, and is of no physical interest.

The transfer matrix T can be computed directly from the $SU(1,1)$ wavefunctions (where care is taken at the singularity [11]):

$$T = \begin{pmatrix} \alpha & \beta \\ \beta^* & \alpha^* \end{pmatrix} \quad (4)$$

$$\begin{aligned} \alpha &= e^{-i\pi m} \left[\frac{\cos \pi m}{\sin \pi(j + \frac{1}{2})} - i \left(\frac{2}{m} \frac{\Gamma(\frac{1-j+m}{2}) \Gamma(\frac{1-m-j}{2})}{\Gamma(\frac{m-1}{2}) \Gamma(-\frac{j+m}{2})} \right. \right. \\ &\quad \left. \left. - \frac{m}{2} \frac{\Gamma(\frac{m+j+1}{2}) \Gamma(\frac{1-m+j}{2})}{\Gamma(\frac{2+m+j}{2}) \Gamma(\frac{2+j-m}{2})} \right) \frac{\cos \pi \frac{j+m}{2} \cos \pi \frac{j-m}{2}}{\sin \pi(j + \frac{1}{2})} \right] \\ \beta &= ie^{i\pi m} \left(\frac{m}{2} \frac{\Gamma(\frac{1+j+m}{2}) \Gamma(\frac{1+j-m}{2})}{\Gamma(\frac{2+j+m}{2}) \Gamma(\frac{2+j-m}{2})} \right. \\ &\quad \left. + \frac{2}{m} \frac{\Gamma(\frac{1-j+m}{2}) \Gamma(\frac{1-m-j}{2})}{\Gamma(-\frac{j-m}{2}) \Gamma(-\frac{j+m}{2})} \right) \frac{\cos \pi \frac{j-m}{2} \cos \pi \frac{j+m}{2}}{\sin \pi(j + \frac{1}{2})}. \end{aligned}$$

Taking $\cos k\pi = \text{Re}(\alpha e^{i\pi m})$, we obtain the dispersion relation $\cos \pi k = \cos \pi m / \sin \pi(j + \frac{1}{2})$, or

$$\mathcal{E}(k) = \frac{1}{\pi^2} \left[\cos^{-1} \left(\cos \pi k \sin \pi \sqrt{A + \frac{1}{4}} \right) \right]^2, \quad (5)$$

which agrees with that derived by Scarf [8]. The bands are restricted to the non-unitary parts of the complementary series of $SU(1,1)$.

Using the properties of our Scarf dynamical symmetry, we can consider a generalization of this potential in $SO(2,2) \approx SU(1,1) \times SU(1,1)$. The states are labeled by j, m, j', c through the direct product $|jm\rangle \times |j'c\rangle$ of $SU(1,1)$ states. It is possible to construct generators of the $SU(1,1)$ algebras, denoted $\mathbf{J}$ and $\mathbf{K}$, for which $J^2 = K^2$, so that $j = j'$, and which has the following periodic Hamiltonian as dynamical symmetry:

$$\begin{aligned} &\left[-\frac{d^2}{dx^2} + \frac{(m+c)^2 - \frac{1}{4}}{\sin^2 x} + \frac{4(j+\frac{1}{2})^2 - \frac{1}{4}}{\cos^2 x} \right] \Psi_{jmc} \\ &= (J_z - K_z)^2 \Psi_{jmc} = (m-c)^2 \Psi_{jmc}. \end{aligned} \quad (6)$$

(This can also be expressed as $V(x) = (a + b \cos x)/\sin^2 x$.) We consider the band structure for $|m+c| \leq \frac{1}{2}$ and $-\frac{3}{4} < j < -\frac{1}{4}$. Again by symmetry, we use the regime $-\frac{1}{2} \leq j < -\frac{1}{4}$. To determine the band structure, we take the projective representations of $SO(2,2)$ obtained through the direct product of the results of Fig. 1 (top). These are shown in Fig. 2, where we plot the complementary series for $SO(2,2)$ as a function of $m-c$ and $m+c$ for three values of j . The shaded areas correspond to unitary representations. The striped

regions are obtained from direct products of unitary representations, while the checkered regions are direct products of non-unitary representations, which can be made unitary in the strip $|m+c| \leq \frac{1}{2}$. The representations in Fig. 2 correspond to (a) $j = -0.45$, (b) $j = -0.35$ and (c) $j = -0.25$. The energy gaps are indicated by the shaded regions. To obtain the bands for a given potential ($m+c$ and j constant), one chooses j and the value of $m+c$ in the range $[-1/2, 1/2]$, and draws the vertical line. The values of $m-c$ which are in the unshaded regions, including the boundaries, correspond to the energy bands. One can see that as $j \rightarrow -\frac{1}{2}$, the bands vanish, leaving only discrete eigenvalues, analogous to Fig. 1(bottom).

The transfer matrix can also be derived (which we omit here for brevity), and results in the dispersion relation:

$$\cos(\pi k) = \frac{\cos(2j+1)\pi \cos(m+c)\pi + \cos(m-c)\pi}{\sin(2j+1)\pi \sin(m+c)\pi}. \quad (7)$$

This $SO(2,2)$ Hamiltonian has three limits where it reduces to the Scarf potential: (a) $j = -\frac{1}{4}$, (b) $m+c = \pm\frac{1}{2}$, and (c) $m+c = 2j+1$. In these cases, the range of $m-c$ agrees with that in Fig. 1 (with twice the period for case (c)), and the dispersion relation agrees with the Scarf result.

We have realized the above dynamical symmetry band problems using projective representations of $SU(1,1)$ in the J^2, J_z basis. There are other ways in which one can realize band structure as well, using bases defined through non-compact generators, such as J^2, J_x . In this case the states $|j, \lambda\rangle$ are continuous, with λ a real number. The final case we would like to discuss here is a more general SGA case of this type. Consider the 1-d crystal formed from the superposition of Pöschl-Teller potentials $V(\rho) = -(j(j+1)/\rho_0^2) \sum_n \cosh^{-2}([\rho - na]/\rho_0)$. Through the linear transformation $\rho = (i\pi\rho_0/2K)x - (a+i\pi\rho_0)/2$, where K and K' are elliptic integrals, the Hamiltonian reduces to the Lamé equation, which has a periodic potential of real period $2K$ [10,12]:

$$\left[-\frac{d^2}{dx^2} + j(j+1)\kappa^2 \text{sn}^2(x|\kappa) \right] \Psi = \mathcal{E} \Psi. \quad (8)$$

Here $\text{sn}(x|\kappa) = \text{sn}(x)$ is the doubly periodic Jacobi elliptic function of modulus κ . This equation can be related to the non-dynamical symmetry problem [5,13]

$$H = J_x^2 + \kappa^2 J_y^2. \quad (9)$$

When the generators $\mathbf{J}$ are written in conical coordinates (θ, ϕ) , the above Hamiltonian decouples into two Lamé equations, one in θ and the other in ϕ . The band edges are obtained from $SU(2)$ [5], which should be contrasted to the previous examples where the edges were classified according to the non-compact algebras $SU(1,1)$ and $SO(2,2)$. For integer j , there are $j+1$ energy bands.

Non-periodic solutions of Lamé's equation can be expressed in terms of theta functions [12]. We can explicitly

construct the transfer matrix and the dispersion relation for the Lamé equation with integer j :

$$\alpha = \cos 2k(\mathcal{E})K \quad \beta = i \frac{\sin 2k(\mathcal{E})K}{\sum_{n=1}^j \text{sn } \alpha_n \text{ dn } \alpha_n / \text{cn } \alpha_n}. \quad (10)$$

where the dispersion relation $k(E)$ is expressed in terms of the Jacobi zeta function Z of modulus κ :

$$k(\mathcal{E}) = -i \sum_{n=1}^j Z(\alpha_n | \kappa^2) + j \frac{\pi}{2K}. \quad (11)$$

The parameters α_n are solutions of j coupled equations which depend upon the eigenvalue $\mathcal{E}$ [12]:

$$\begin{aligned} \mathcal{E} &= \sum_{n=1}^j \text{sn}^2 \alpha_n - \left[\sum_{n=1}^j \text{cn } \alpha_n \text{ ds } \alpha_n \right]^2 \\ 0 &= \sum_{p=1}^j \frac{\text{sn } \alpha_p \text{ dn } \alpha_p \text{ cn } \alpha_p - \text{sn } \alpha_n \text{ dn } \alpha_n \text{ cn } \alpha_n}{\text{sn}^2 \alpha_p - \text{sn}^2 \alpha_n} \quad (p \neq n). \end{aligned} \quad (12)$$

This transfer matrix has the following properties. (a) In the limit $\kappa = 1$, the Hamiltonian becomes the Pöschl-Teller potential, which is not periodic, and the denominator of β becomes $i\sqrt{\mathcal{E} - j(j+1)} = ik$. Here we obtain the transfer matrix with matrix elements $\alpha_{PT} = \Gamma(ik)\Gamma(1+ik)/\Gamma(1+ik+j)\Gamma(ik-j)$ and $\beta_{PT} \rightarrow 0$, the desired result for integer j . (b) In the limit $\kappa = 0$, the Hamiltonian (9) becomes $H = J_x^2$ while (8) is free motion. In the basis $|j, \lambda\rangle$, (9) has the desired free particle dispersion, $\mathcal{E} = \lambda^2 \geq 0$. We also recover the free particle transfer matrix. (c) There are additional connections one can make with the Mathieu equation when $\kappa \rightarrow 0$, $j \rightarrow \infty$, and $\kappa^2 j(j+1) = \text{const.}$, as well as connections to the Scarf potential. Because the equations (12) must generally be solved numerically, we show the type of result one finds for $j = 1$. Here there is one root α , and the wavefunction in Bloch form is:

$$\Psi_k(x) = \left[\frac{H(x+\alpha)}{\Theta(x)} \exp\left(\frac{i\pi x}{2K}\right) \right] \exp(-ikx), \quad (13)$$

where H and Θ are theta functions, and α is determined from $\mathcal{E} = \text{dn}^2 \alpha + \kappa^2$, which restricts the energy to the bands $\kappa^2 \leq \mathcal{E} \leq 1$ and $1 + \kappa^2 \leq \mathcal{E} < \infty$. For the lower band, $\alpha = K + i\beta$ where β ranges from K' to 0, while for the upper band $\alpha = i\beta$, where β ranges from 0 to K' . The dispersion relation then has the form

$$k(\mathcal{E}) = -Z(\beta|1-\kappa^2) + \frac{\pi}{2K} \left(1 - \frac{\beta}{K'} \right) + \mathcal{E}', \quad (14)$$

where $\mathcal{E}' = \sqrt{(1-\mathcal{E})(\mathcal{E}-\kappa^2)/(1+\kappa^2-\mathcal{E})}$ for the lower band, and $\mathcal{E}' = \sqrt{(\mathcal{E}-\kappa^2)(\mathcal{E}-1-\kappa^2)/(\mathcal{E}-1)}$ for the upper band. In contrast to the Scarf problem, we see here that the band edges are given by the discrete representations of $SU(2)$. The bands correspond to a diagonalization in the $SU(1,1)$ basis, such as that defined by

J and J_x . This is evident if we examine the limit $\kappa = 0$, where we expect a continuous spectrum.

We have shown that SGA and dynamical symmetry techniques can be applied to Hamiltonians with periodic potentials, and band structure can arise naturally from representation theory. This fills a long-standing gap in the algebraic approach to quantum systems. Hamiltonians such as the Scarf's are now reduced to problems as simple as solving $H = J_z^2$, using representations with multi-valued wavefunctions. The complementary series now finds natural application in the band structure problem, where its non-unitary representations are important. Further, both compact and non-compact Lie algebras play a role in the band edges - $SU(1,1)$ for the Scarf and extended Scarf potentials, and $SU(2)$ for the Lamé equation. We expect that dynamical symmetries exist for potentials in higher dimensions, where the point group of the lattice will play a role in the representations of algebras such as $SU(n, m)$. Within these representations one should find the complete spectral solutions to the problems.

We would like to thank F. Iachello for many useful discussions. This work was supported by DOE grant DE-FG02-91ER40608.

ysis (Cambridge University Press, London, 1963).

[13] J. Patera, P. Winternitz, *J. Math. Phys.* **14** (1973) 1130.




FIG. 1. (Top) Complementary series of $SU(1,1)$ labeled by j, m . The unitary (projective) representations satisfy $m_0(m_0 - 1) > j(j + 1)$, where $m = m_0 \pm n$ ($n = 0, 1, \dots$) and $0 \leq m_0 < 1$, and coincides with the energy gaps. (Bottom) The corresponding Scarf energy spectrum $E = [\hbar^2/(2M\rho_0^2)]m^2$ is plotted as a function of j . The band edges are given by the discrete series and the bands (shaded) by the non-unitary part of the complementary series.

- [1] Y. Dothan, M. Gell-Mann and Y. Ne'eman, *Phys. Lett.* **17** (1965) 148; F. Gürsey and L. Radicati, *Phys. Rev. Lett.* **13** (1964) 173; A. O. Barut and A. Bohm, *Phys. Rev.* **B139** (1965) 1107. For a historical review, see also F. Iachello, *Rev. Nuovo Cimento* **19** (1996) 1.
- [2] F. Iachello and A. Arima, *The Interacting Boson Model* (Cambridge Press, Cambridge, 1987).
- [3] F. Iachello and R. Levine, *Algebraic Theory of Molecules* (Oxford Press, Oxford, 1995).
- [4] D. Kusnezov, *Phys. Rev. Lett.* **79** (1997) 537.
- [5] Y. Alhassid, F. Gursey and F. Iachello, *Phys. Rev. Lett.* **50** (1983) 873.
- [6] A. Frank, K. B. Wolf, *Phys. Rev. Lett.* **52** (1984) 1737.
- [7] Y. Alhassid, F. Gürsey, F. Iachello, *Ann. Phys. (NY)* **148**, 346 (1983); *Ann. Phys. (NY)* **167**, 181 (1983).
- [8] F. L. Scarf, *Phys. Rev.* **112**, 1137 (1958); A. M. Perelomov, *Sov. J. Part. Nucl.* **10**, 336 (1979); L.D. Salem, R. Montemayor, *Phys. Rev.* **A47** 105, (1993); R. De, R. Dutt, U. Sukhatme, *J. Phys. A: Math. Gen.* **25** L843 (1992); F. Gesztesy, C. Macdeo, L. Streit, *J. Phys. A: Math. Gen.* **18** L503 (1985).
- [9] V. Bargmann, *Ann. Math.* **48**, 568 (1947); L. Pukánszky, *Math. Annalen* **156**, 96 (1964).
- [10] B. Sutherland, *Phys. Rev.* **A 8** (1973) 2514.
- [11] H. Li, D. Kusnezov, in *Group22 Proceedings*, Eds. S. P. Corney, R. Delbourgo, P.D. Jarvis, (International, Cambridge, MA, 1999), p. 310.
- [12] E.T. Whittaker, G.N. Watson, *A Course in Modern Anal-*




FIG. 2. Unitary representations (shaded regions) of $SO(2,2)$ obtained from Fig. 1 (top), restricted to the physical region $|m + c| \leq 1/2$, for selected values (a) $j = -0.45$, (b) $j = -0.35$ and (c) $j = -0.25$. The physical spectrum, given by $(m - c)^2$, is obtained by choosing a value of $m + c$ and j (i.e. a specific potential). The bands are the values of $m - c$ which are non-unitary (in the unshaded regions).